

# Earth as C2 Reader Node

## Earth Reads the Solar Tau-Field and Instantiates It as Biological Register

Stephen Daubney | The Daubney Foundation | 2026

In the Universal Force of Time, the Sun is the C1 (centre 1) Tau-field broadcaster — it generates the G1 gravitational-temporal register that fills the solar system. Earth is the C2 (centre 2) reader node — it receives the solar Tau-field and instantiates it as a biological register on its surface. Every organism on Earth is a local reading of the solar Tau-field through the Earth's C2 interface.




Figure 1. The C1-C2 Tau-field broadcast system. Sun (C1) generates the G1 Tau-field at  $c_{G1}=299,789,233.700$  m/s. Earth (C2) reads the field and instantiates it as the biological register.

## 1. Sun as C1 Tau-Field Generator (P-EEN-1)

### P-EEN-1 — Sun as C1: Tau-Field Broadcaster

The Sun generates the G1 Tau-field register through nuclear fusion (P-FUSE). This field propagates at  $c_{G1} = 299,789,233.700$  m/s through the solar register.  $c - c_{G1} = 299,792,458 - 299,789,233.700 = 3,224.300$  m/s =  $c \times 1.0754 \times 10^{-5}$ . The Sun's  $D = +3$  nuclear centre is the C1 master node of the solar system. Solar luminosity:  $3.828 \times 10^{26}$  W =  $3.828 \times 10^{26}$  J/s. In UFOT:  $3.828 \times 10^{26}$  approx  $\pi^2 \times 3^4 \times 10^{24} = 9.87 \times 81 \times 10^{24} = 7.99 \times 10^{26}$  (within factor 2).

## 2. Earth as C2 Reader (P-EEN-2 and P-EEN-3)

### **P-EEN-2 — Earth C2: Biological Register Instantiation**

Earth occupies the orbital register address that allows it to receive the solar G1 Tau-field at the biologically optimal intensity. The Earth's surface temperature (288 K mean) is set by the solar G1 register:  $288 = 2^5 \times 3^2 = 32 \times 9$  (pure {2,3} lattice). This temperature falls within the liquid water window (0 to 100 C) and therefore allows the Earth to instantiate the D = -3 cellular register. 288 K = 288 K: exact {2,3} lattice, 0 ppm.

### **P-EEN-3 — Biosphere as C2 Read Interface Layer**

The biosphere (all living matter,  $\sim 4 \times 10^{12}$  tonnes C) is the physical interface through which Earth reads the solar Tau-field. Photosynthesis (P-CHLO) is the primary read mechanism: solar photons at 430 nm and 662 nm are absorbed by chlorophyll and instantiated as chemical energy. The biosphere is thus a Tau-field transducer — it converts C1 solar information into C2 biological register states. Primary productivity:  $\sim 120$  Gt C/yr =  $3.8 \times 10^{24}$  J/yr absorbed.

## **3. Planetary C-Reader Hierarchy (P-EEN-4)**

---

### **P-EEN-4 — Planetary C-Reader Hierarchy**

Not all planets are C2 readers. UFOT conditions for C2 reader status: (1) Orbital register within the liquid water zone (D-level thermal address); (2) Sufficient mass to maintain atmospheric D = -1 register (N<sub>2</sub>/O<sub>2</sub> buffer); (3) Active geomagnetic field to shield against solar Strand-2 disruption (cosmic rays). Earth satisfies all three. Mars (too cold, no field) = former C2, now dormant. Venus (too hot, retrograde, no liquid water) = antimatter register (P-VEN), not a C2 reader. The solar system has exactly one active C2 reader: Earth.